q

arXiv:math/9706203v1 [math.GR] 4 Jun 1997

ROTATING RATIONALLY

by
Charles Radin[1] and Lorenzo Sadun[2]

Mathematics Department
University of Texas
Austin, TX 78712

radin@math.utexas.edu and sadun@math.utexas.edu

---

1 Research supported in part by NSF Grant No. DMS-9531584
2 Research supported in part by NSF Grant No. DMS-9626698

§1. Introduction

In this paper we will be considering algebraic features of $SO(3)$, the group of rotations of three dimensional Euclidean space. One can view $SO(3)$ as the ways in which a solid object, such as a prism, is oriented in space. Figure 1 shows a portion of the "quaquaversal" tiling [CR] of Euclidean space by prisms. In this tiling the prisms seem to be pointing in all directions! Indeed, prisms in this tiling appear in an infinite number of orientations, and these orientations form a dense subgroup of $SO(3)$. Amazingly, this subgroup (which we denote $G(6,4)$) is generated by only two elements, a rotation by $\pi/3$ about the x axis and a rotation by $\pi/2$ about the z axis.

We generalize this situation as follows. Let A and B be finite order rotations (i.e. rotations by rational multiples of π), about axes that are themselves separated by an angle of finite order. We are interested in understanding the algebraic relations between A and B. One consequence of our analysis is a classification of all subgroups of $SO(3)$ generated by such a pair of rotations. We will show that all such relations can be understood in terms of a short list of simple relations involving rotations by $\pi/2$:

1. If A is a rotation by π about an axis (say, the x axis), and B is any rotation about an orthogonal axis (say, the z axis), then $ABA = B^{-1}$. Equivalently, $ABAB$ is the identity.
2. If A and B are rotations by $\pi/2$ about orthogonal axes, then $ABABAB$ is the identity. Equivalently, $ABA = BAB$. (For example, if A is a positive rotation about the y axis and B is a positive rotation about the z axis, then AB cyclically permutes the positive x, y, and z axes.)
3. If A and B are rotations by π about axes that are separated by an angle θ, then BA is a rotation, by 2θ, about an axis perpendicular to the axes of A and B. (For example, if A is a rotation about the x axis, and the axis of B lies in the x-y plane, then BA is a rotation by 2θ about the z-axis.

Before we begin our analysis, we must explain just what we mean by a "relation" between two elements of a group. That requires a digression into the language of group presentations.

Let A be a nonempty collection of pairs of symbols a_j, a_j^{-1}. We call each symbol a_j a "generator". We define "words" to be finite ordered sequences of such symbols, such as $a_3 a_1^{-1} a_{17}$. The empty ordered sequence is denoted 1. A product is defined for such sequences by concatenation, for instance the product $(a_3 a_1^{-1} a_{17})(a_{12} a_4) = a_3 a_1^{-1} a_{17} a_{12} a_4$. The symbol a^n is an abbreviation for the word $aaa \cdots a$ consisting of n a's. Given a word $W = abc \cdots jk$, the word W^{-1} is defined to be $W^{-1} = k^{-1} j^{-1} \cdots c^{-1} b^{-1} a^{-1}$.

Assume given, besides A, some (possibly empty) set of words $P_1, P_2, \cdots$, which we call relators. These relators are the words that we wish to set equal to the identity. We do this formally by defining an equivalence relation on the set of words.

Let $W_1 \sim W_2$ if W_1 can be turned into W_2 by a finite number of applications of the following operations:

- Insertion of one of the words $a_j a_j^{-1}$, $a_j^{-1} a_j$, P_j, or P_j^{-1}, either between two consecutive symbols in W_1, or just before the first symbol of W_1 or just after the last symbol of W_1.

- Deletion of one of the words $a_j a_j^{-1}$, $a_j^{-1} a_j$, P_j, or P_j^{-1}, if it forms a consecutive block of symbols in W_1.

We denote the set of equivalence classes of words by $< a_1, a_2, \cdots : P_1, P_2, \cdots >$. It is easily seen to be a group with product defined by concatenation, and the expression $< a_1, a_2, \cdots : P_1, P_2, \cdots >$ is called a "presentation" of the group. The equations $P_i = 1$ are called "relations", as are all equations that can be derived from the basic relations $P_i = 1$.

In this notation a presentation of the cyclic group of order n, $\mathbb{Z}_n$, is given by $< a_1 : a_1^n >$, and a presentation of the dihedral group of order $2n$, D_n, is given by $< a_1, a_2 : a_1^n, a_2^2, (a_1 a_2)^2 >$. For another example consider the two finite cyclic groups $\mathbb{Z}_3$, $\mathbb{Z}_5$, of orders 3 and 5 respectively. Their "free product", denoted $\mathbb{Z}_3 * \mathbb{Z}_5$, is the group $< a_1, a_2 : a_1^3, a_2^5 >$; that is, all finite strings of a_1's and a_2's of appropriate powers.

There is a generalization of free products which is of significant interest. Consider the group G with presentation $< a, b : a^4, b^6, a^2 b^{-3} >$. This is almost the free product of $\mathbb{Z}_4 = < a : a^4 >$ with $\mathbb{Z}_6 = < b : b^6 >$. Both of these groups have $\mathbb{Z}_2$ as a subgroup — the subgroups generated by a^2 and b^3 respectively — and the relator $a^2 b^3$ identifies these two subgroups as the same. G is said to be the *amalgamated free product* of the subgroups generated by a and b, respectively, over the common subgroup $\mathbb{Z}_2$. (Think of this as an algebraic version of the process in topology whereby you "identify" the opposite edges of a square to get a torus.) This is expressed $G = \mathbb{Z}_4 *_{\mathbb{Z}_2} \mathbb{Z}_6$, although this notation does not show the way in which the common subgroups of the factors are identified. More generally, the group $G \equiv < a_1, a_2, \cdots, b_1, b_2, \cdots, : R_1(\{a_i\}), \cdots, S_1(\{b_j\}), \cdots, U_1(\{a_i\}) V_1^{-1}(\{b_j\}), \cdots >$ is called the "amalgamated free product" of the subgroup A, generated by the a's with relations $\{R_i\}$, and the subgroup B, generated by the b's with relations $\{S_j\}$, in which the subgroup of A generated by the U_i's is identified with the subgroup of B generated by the V_j's. If this common subgroup is H, we write $G = A *_H B$.

We mentioned above that we will classify all subgroups of $SO(3)$ generated by a pair of rotations of finite order about axes that are themselves separated by an angle of finite order. For each such group we will give an explicit presentation. In many cases we can also express the group as a free product, or an amalgamated free product, of simpler groups.

We need some further notation. See Figure 2. Let ℓ be the line in the x-y plane, through the origin and making an angle of $2\pi n/m$ with the x-axis, where n and m are relatively prime positive integers and $m > 2$. Let $A = R_x^{2\pi/p}$ be rotation by

$2\pi/p$ about the x-axis, where p is a positive integer. Let $B = R_\ell^{2\pi/q}$ be rotation by $2\pi/q$ about the line ℓ. A and B will be our primary objects of study. We similarly define the rotations $C = R_z^{\pi/q}$, $S = R_y^{\pi/2}$ and $U = R_z^{2\pi/m}$. T and $\hat{T}$ will denote various rotations about the x axis.

We denote the subgroup of $SO(3)$ generated by A and B as $G_{n/m}(p,q)$. An important special case is where the axes of A and B are orthogonal: that is, $n/m = 1/4$. In that case we omit the subscript and write $\hat{G}(p,q) \equiv G_{1/4}(p,q)$. Among the groups $\hat{G}(p,q)$, a further special case is where $q = 4$, which will play a pivotal role.

These groups form a hierarchy. The structure of $\hat{G}(p,4)$ is considerably simpler than that of a general $\hat{G}(p,q)$, while the structure of $\hat{G}(p,q)$ is much simpler than that of a general $G_{n/m}(p,q)$. However, $G_{n/m}(p,q)$ actually embeds in the "simpler" $\hat{G}(m,pq)$, while $\hat{G}(p,q)$ embeds in the "simpler" $\hat{G}(pq,4)$:

$$G_{n/m}(p,q) \subset \hat{G}(p',q') \subset \hat{G}(s,4) = \text{ not too complicated.} \tag{1}$$

We can therefore use the known properties of a simpler group to deduce the structure of the more complicated group! The cases $\hat{G}(3,3)$ and $\hat{G}(3,4)$ were worked out in [CR]; the structures of $\hat{G}(p,4)$ and $\hat{G}(p,q)$ were derived in [RS]; the structure of $G_{n/m}(p,q)$ is new.

To see the embeddings of (1) it is useful to define $G(p,q)$ as the subgroup (conjugate to $\hat{G}(p,q)$) generated by rotations about the x and z-axes, rather than the x and y-axes. That is, $G(p,q)$ is generated by A and C, rather than A and B, while $\hat{G}(m,4)$ is generated by $T = R_x^{2\pi/m}$ and $S = R_y^{\pi/2}$. Note that, if s is a multiple of p and also a multiple of q, then $A = T^{s/p}$, while $C = S^{-1}T^{s/q}S$, so that $G(p,q) \subset \hat{G}(s,4)$. Next we show that $G_{n/m}(p,q) \subset G(r,m)$, where r is any common multiple of p and q. The generators of $G_{n/m}(p,q)$ are $A = R_x^{2\pi/p}$ and $B = R_\ell^{2\pi/q}$, while the generators of $G(r,m)$ are $\hat{T} = R_x^{2\pi/r}$ and $U = R_z^{2\pi/m}$. Since r is a multiple of p and q, $A = \hat{T}^{r/p}$ and $B = U^{-n}\hat{T}^{r/q}U^n$.

Although there are quite a few cases, depending on the values of p, q, m and n, the results of this paper, and of [RS], are all of the following form:

"Theorem" *Let* $G = \hat{G}(m,4)$, $G(p,q)$ *or* $G_{n/m}(p,q)$. *There is a short list of simple relations among the generators of* G*, all of which involve rotations by multiples of* $\pi/2$*, and all of which are simple consequences of*

$$R_x^\pi R_y^\theta R_x^\pi = R_y^{-\theta}, \tag{2}$$

$$R_y^{\pi/2} R_x^{\pi/2} R_y^{\pi/2} = R_x^{\pi/2} R_y^{\pi/2} R_x^{\pi/2}, \tag{3}$$

$$R_\ell^\pi R_x^\pi = R_z^{4\pi n/m}. \tag{4}$$

Any other relation may be derived from these simple relations.

For example, powers of the generators of $G_{n/m}(p,q)$, with p and q both odd, are never rotations by multiples of $\pi/2$, unless they are rotations by 2π. So we expect (and prove) that, if p and q are odd, $G_{n/m}(p,q)$ is the free product of $\mathbb{Z}_p$ and $\mathbb{Z}_q$, for any n and m.

More precise formulations of this "theorem" may be found below. The cases $G = \hat{G}(m,4)$ and $G(p,q)$ are discussed in §2 (Theorems 1 and 2). The cases $G_{n/m}(p,q)$ fall into 3 classes, which are the subjects of Theorems 3–5 in §3. Theorems 3–5 depend on Theorem 2, which in turn depends on Theorem 1, which in turn depends on the following result:

Foundation Lemma: *Let* $T = R_x^{2\pi/m}$, *let* $S = R_y^{\pi/2}$, *and consider the word*

$$\chi = S^W S^{a_1} T^{b_1} S^{a_2} T^{b_2} \cdots S^{a_n} T^{b_n} S^E, \tag{5}$$

where W, E, a_i, b_i *and* n *are integers,* $n > 0$, *none of the* a_i*'s are divisible by 2, and none of the* b_i*'s are divisible by* $m/4$. *Then* χ *is not equal to the identity.*

The proof of the Foundation Lemma may be found in [RS]; here is a sketch. We write S and T as explicit 3×3 real matrices. Using commutative ring theory, we show that the four upper right hand matrix elements of $\Pi_i(S^{a_i}T^{b_i})$ are not integers, and hence χ has four non-integral matrix elements, and hence χ is not the identity matrix.

The following two extensions of the Foundation Lemma will also be used repeatedly:

Extension 1: *Let* $T = R_x^{2\pi/m}$, *let* $S = R_y^{\pi/2}$, *and let* χ *be as in eq (5). If* W, E, a_i, b_i *and* n *are integers,* $n > 0$, *none of the* a_i*'s are divisible by 2, and no two* consecutive b_i*'s are divisible by* $m/4$, *then* χ *is not equal to the identity.*

Proof: If $b_i = m/4$, $a_{i-1} = a_i = 1$ and b_{i-1} and b_{i+1} are not divisible by $m/4$, then we can shorten our word, removing the $T^{m/4}$ term using the identity (3):

$$T^{b_{i-1}} S T^{m/4} S T^{b_{i+1}} = T^{b_{i-1}} T^{m/4} S T^{m/4} T^{b_{i+1}} = T^{b_{i-1}+m/4} S T^{b_{i+1}+m/4}. \tag{6}$$

Similar manipulations apply if a_{i-1}, b_i or a_i is negative. Note that, since $b_{i\pm1}$ is not divisible by $m/4$, neither is $b_{i\pm1} + m/4$. We repeat this procedure until all the factors of $T^{\pm m/4}$ are removed, except possibly T^{b_1} and T^{b_n}. If b_1 and b_n are not multiples of $m/4$, the Foundation Lemma applies, and we are done. If b_1 or b_n equals $\pm m/4$ we apply the proof of the Foundation Lemma to the terms in the middle (e.g., $S^{a_2}T^{b_2} \cdots S^{a_n}$ if $b_1 = b_n = m/4$) and see that this central piece has four non-integral matrix elements. Since S and $T^{m/4}$ are, up to sign, permutation matrices, χ has four non-integral matrix elements and cannot equal the identity. ∎

Extension 2: *Let* A *and* C *be rotations about orthogonal axes by* $2\pi/p$ *and* $2\pi/q$, *respectively, and consider a word of the form*

$$A^{a_1} C^{c_1} \cdots A^{a_n} C^{c_n}, \tag{7}$$

where a_1 *and* c_n *may be zero, but otherwise none of the* a_i*'s are multiples of* $p/2$ *and none of the* c_j*'s are multiples of* $q/2$. *If no two consecutive terms represent* *rotations by* $\pm\pi/2$ *(e.g* $A^{p/4}C^{-q/4}$ *or* $C^{q/4}A^{p/4}$*), and if the word is not empty, then the word is not equal to the identity.*

Proof: Choose axes such that $A = R_x^{2\pi/p}$ and $C = R_z^{2\pi/q}$, and let $S = R_y^{\pi/2}$ and $T = R_x^{2\pi/pq}$, so that $A = T^q$ and $B = S^{-1}T^pS$. Rewrite the word in terms of S and T and apply Extension 1. ∎

To prove Theorems 1–5, we employ two strategies repeatedly. We always begin by noting certain simple relations among the generators of G. The first strategy is to use these relations to convert an arbitrary word in the generators either into the identity or into a word which, by the Foundation Lemma and its Extensions, is known not to equal the identity. When this strategy succeeds it implies that there cannot be any relations that are independent of the stated ones. A presentation of the group then follows. A second strategy is to embed the group G in a seemingly larger group H, and then use the simple relations to show that the generators of H are actually in G, and thus that $G = H$. All cases yield to a combination of these strategies. In a small number of cases (Theorem 5), both strategies are needed. First we choose an appropriate H and show that $G = H$, and then we apply the first argument to H to find a presentation.

§2. The structures of $\hat{G}(m, 4)$ and $G(p, q)$.

Before examining the structures of $G(p, q)$ and $\hat{G}(m, 4)$ in general, we consider the few cases where the group is finite. $G(p, 1)$ is the cyclic group of order p, and has the presentation

$$G(p,1) = \mathbb{Z}_p = <\alpha : \alpha^p> . \tag{8}$$

$G(p, 2)$ is the dihedral group of order $2p$, and has the presentation

$$G(p,2) = D_p = <\alpha, \beta : \alpha^p, \beta^2, (\alpha\beta)^2> . \tag{9}$$

The relation $(\alpha\beta)^2 = 1$, which is equivalent to eq. (2), will be used repeatedly.

The last finite group is $\hat{G}(4, 4) = G(4, 4)$, which is the 24-element group of rotational symmetries of the cube. It has the presentation

$$\hat{G}(4,4) = <\alpha, \beta : \alpha^4, \beta^4, (\alpha\beta^2)^2, (\alpha^2\beta)^2, (\alpha\beta)^3> . \tag{10}$$

A new ingredient is the last relator in (10), which is a consequence of eq. (3). This, too, will be used frequently.

Theorem 1: *The group* $\hat{G}(m, 4)$ *has the following presentation:*

1. *If* m *is odd, then*

$$\hat{G}(m,4) = <\alpha, \beta : \alpha^m, \beta^4, (\alpha\beta^2)^2> = D_m *_{\mathbb{Z}_2} \mathbb{Z}_4, \tag{11}$$

2. *If* m *is twice an odd number, then*

$$\hat{G}(m,4) = <\alpha, \beta : \alpha^m, \beta^4, (\alpha\beta^2)^2, (\alpha^{m/2}\beta)^2> = D_m *_{D_2} D_4, \tag{12}$$

3. *If m is divisible by 4, then*

$$\hat{G}(m,4) = <\alpha,\beta : \alpha^m, \beta^4, (\alpha\beta^2)^2, (\alpha^{m/2}\beta)^2, (\alpha^{m/4}\beta)^3> = D_m *_{D_4} \hat{G}(4,4). \tag{13}$$

Proof: Let $\hat{G}(m,4)$ be generated by $S = R_y^{\pi/2}$ and $T = R_x^{2\pi/m}$. In each case, the relators listed (with α corresponding to T and β to S), are just the previously discussed relators either involving rotations by π or involving pairs of rotations by $\pi/2$. Our task is to show there are no additional relations. We do this by taking an arbitrary word in S and T and, using the known relations, either reducing the word to the form (5) or to the identity. If the reduction is to the form (5), then, by the Foundation Lemma (or Extension 1), the word cannot be a relator. If the reduction is to the identity, the word *is* a relator, but is not independent of the listed relators.

We begin with case 1. An arbitrary word in S and T is of the form

$$S^W S^{a_1} T^{b_1} S^{a_2} T^{b_2} \cdots S^{a_n} T^{b_n} S^E, \tag{14}$$

with W, a_i, b_i and E arbitrary. Using the relators S^4 and T^m, we can require that the b_i's be between 1 and $m-1$, inclusive, and that each a_i be 1, 2 or 3. Since m is odd, T^{b_i} can never be a rotation by a multiple of $\pi/2$. If any of the a_i's equal 2, we can shorten the word using $(TS^2)^2 = 1$, or equivalently $T^b S^2 = S^2 T^{-b}$, and hence $T^b S^2 T^{b'} = S^2 T^{b'-b}$. In this way, we shorten the word until either $n = 0$ or until all the a_i's are odd, in which case we have the form (5).

In case 2, T^{b_i} may be a rotation by π, but not by $\pm\pi/2$. In that case, in addition to removing S^2 factors, we remove $T^{m/2}$ factors, using $T^{m/2}S = S^{-1}T^{m/2}$ and hence $T^{m/2}S^a = S^{-a}T^{m/2}$. Once the $T^{m/2}$s and S^2s are removed, we have the form (5).

In case 3, even after removing the $T^{m/2}$s and S^2s, we may have some $T^{\pm m/4}$s in our word. If two consecutive factors appear we shorten the word using $(T^{m/4}S)^3 = 1$, and hence $ST^{m/4}ST^{m/4} = T^{-m/4}S^{-1}$, with similar identities applying if the exponents are not all positive. Thus the word may continue to be shortened until either Extension 1 may be applied, or the word is so short it can be checked by hand.

This gives the presentations (11–13). As for the amalgamated free products, in case 1 the D_m subgroup is generated by α and β^2, $\mathbb{Z}_4$ is generated by β, and their intersection is $\mathbb{Z}_2 = \{1, \beta^2\}$. In case 2 the D_m subgroup is generated by α and β^2, the D_4 subgroup by $\alpha^{m/2}$ and β, and the common D_2 by $\alpha^{m/2}$ and β^2. In case 3, D_m is generated by α and β^2, $\hat{G}(4,4)$ by $\alpha^{m/4}$ and β, and the common D_4 subgroup by $\alpha^{m/4}$ and β^2. ■

Theorem 2: *The group $G(p,q)$ has the following structure:*

1. *If p and q are both odd, then*

$$G(p,q) = <\alpha,\beta : \alpha^p, \beta^q> = \mathbb{Z}_p * \mathbb{Z}_q, \tag{15}$$

2. If p is even and q is odd, then

$$G(p,q) = <\alpha, \beta : \alpha^p, \beta^q, (\alpha^{p/2}\beta)^2> = \mathbb{Z}_p *_{\mathbb{Z}_2} D_q, \tag{16}$$

3. If p and q are both even, but q is not divisible by 4, then

$$G(p,q) = <\alpha, \beta : \alpha^p, \beta^q, (\alpha^{p/2}\beta)^2, (\alpha\beta^{q/2})^2> = D_p *_{D_2} D_q. \tag{17}$$

4. If p and q are both divisible by 4, then $G(p,q) = \hat{G}([p,q],4)$.

Proof: In cases 1–3, the relations listed are known consequences of rotations by π. What remains is to show that these are the only relations.

Let $A = R_x^{2\pi/p}$ and $C = R_z^{2\pi/q}$ be the generators of $G(p,q)$. Take an arbitrary word in A and C, and use the given relations as above to put it in the form

$$A^{a_1}C^{c_1}\cdots A^{a_n}C^{c_n}, \tag{18}$$

where a_1 and c_n may be zero, but none of the other a_i's are multiples of $p/2$, and none of the other c_j's are multiples of $q/2$. (In case 1 there is nothing to do, in case 2 we are eliminating factors of $A^{p/2}$, and in case 3 we are eliminating factors of $A^{p/2}$ and $C^{q/2}$). Since q is not divisible by 4, none of the remaining C^{c_j} terms are rotations by multiples of $\pi/2$. Thus, by Extension 2, our word is not the identity.

All that remains for cases 1–3 are the amalgamated free products. In case 2, D_q is generated by $A^{p/2}$ and C, while $\mathbb{Z}_p$ is generated by A. In case 3, D_p is generated by A and $C^{q/2}$, D_q is generated by $A^{p/2}$ and C, and their intersection, D_2, is generated by $A^{p/2}$ and $C^{q/2}$.

In case 4, we consider S and $\hat{T} = R_x^{2\pi/[p,q]}$, the generators of $\hat{G}([p,q],4)$. Since A and C can be constructed from S and $\hat{T}$, we must have $G(p,q) \subseteq \hat{G}([p,q],4)$. We will show that $S, \hat{T} \in G(p,q)$, and thus $G(p,q) = \hat{G}([p,q],4)$.

Since p is divisible by 4, $R_x^{\pi/2} = A^{p/4} \in G(p,q)$. Since q is divisible by 4, $R_z^{\pi/2} = C^{q/4} \in G(p,q)$. Thus $S = R_y^{\pi/2} = R_x^{-\pi/2}R_z^{\pi/2}R_x^{\pi/2} \in G(p,q)$. Now $S^{-1}CS = R_x^{2\pi/q} \in G(p,q)$. Since $R_x^{2\pi/p}$ and $R_x^{2\pi/q}$ are in the group, so is $\hat{T}$. ■

§3. The structure of $G_{n/m}(p,q)$.

In studying $G(p,q)$ (Theorem 2) we saw that one of two things happened: either the only relations among the generators A and C were the obvious ones, or $G(p,q)$ was equal to the previously understood group $\hat{G}([p,q],4)$. In analyzing $G_{n/m}(p,q)$, we shall have 3 possibilities. Sometimes there are no non-trivial relations between the generators; these cases are treated in Theorem 3. Sometimes $G_{n/m}(p,q) = G(m,[p,q])$; these cases are treated in Theorem 4. In a few exceptional cases (Theorem 5), $G_{n/m}(p,q)$ is neither $\mathbb{Z}_p * \mathbb{Z}_q$ nor $G(m,[p,q])$; these cases are in some sense intermediate.

Our notation is as in Figure 2. As always, $A = R_x^{2\pi/p}$, $S = R_y^{\pi/2}$, ℓ is the line in the x-y plane, through the origin, making an angle of $2\pi n/m$ with the x axis, and $B = R_\ell^{2\pi/q}$. We now fix $T = R_x^{2\pi/pq}$, $\hat{T} = R_x^{2\pi/[p,q]}$ and $U = R_z^{2\pi/m}$. A and B generate $G_{n/m}(p,q)$, while U and T (or $\hat{T}$) generate $G(pq,m)$ (or $G([p,q],m)$). Note that $A = T^q$ and $B = U^{-n}T^pU^n$, so $G_{n/m}(p,q) \subseteq G(m,pq)$. Similarly, $G_{n/m}(p,q) \subseteq G([p,q],m)$. For any integer N, let $\rho(N)$ be the number of powers of 2 that divide N. For example $\rho(3) = 0$ and $\rho(12) = 2$.

Without loss of generality we can assume that m is either odd or a multiple of 4. For if m is an odd multiple of 2, then we can replace the axis ℓ by $-\ell$, which makes an angle of $2\pi([(m/2)+n]/m)$ with the x axis. Since $m/2$ and n are odd, $m/2+n$ is even, and we can replace n and m by $n' = (m/2+n)/2$ and $m' = m/2$, where m' is odd.

Theorem 3: *If p and q are odd, or if p is even, q is odd and $m \neq 4$, then*

$$G_{n/m}(p,q) = <\alpha, \beta : \alpha^p, \beta^q> = \mathbb{Z}_p * \mathbb{Z}_q. \tag{19}$$

If p is even, q is odd and $m = 4$, then

$$G_{n/m}(p,q) = <\alpha, \beta : \alpha^p, \beta^q, (\alpha^{p/2}\beta)^2> = \mathbb{Z}_p *_{\mathbb{Z}_2} D_q. \tag{20}$$

Proof: Take an arbitrary nontrivial word in A and B,

$$A^{a_1}B^{b_1} \cdots A^{a_n}B^{b_n}, \tag{21}$$

and rewrite it in terms of U and T:

$$T^{qa_1}U^{-n}T^{pb_1}U^n \cdots T^{pb_n}U^n. \tag{22}$$

If p and q are odd then no power of T can be a rotation by $\pi/2$ or π. Since $m > 2$, and since n and m are relatively prime, $U^{\pm n}$ is not a rotation by π (although it may be a rotation by $\pi/2$ if $m = 4$). By Extension 2 the word is not equal to the identity, so no relations between A and B exist.

If p is even, q is odd, and $m \neq 4$, then U^n is not a rotation by $\pi/2$ or π. However, some of the T^{qa_i} terms may be rotations by π. We remove these using the fact that

$$T^{pb_{i-1}}U^nT^{pq/2}U^{-n}T^{pb_i} = T^{pb_{i-1}}U^{2n}T^{pb_i+pq/2}. \tag{23}$$

Note that $T^{pb_i+pq/2}$ is not a rotation by a multiple of $\pi/2$. There may remain some T^{qa_i} terms that are rotations by $\pi/2$, and U^{2n} might be a rotation by $\pi/2$ (if $m = 8$), but these cannot occur consecutively. Every T^{qa_i} is flanked by $U^{\pm n}$s, while every U^{2n} is flanked by $T^{pb_{i-1}}$ and $T^{pb_i+pq/2}$. Thus by Extension 2 the word is not equal to the identity.

The case where p is even, q is odd and $m = 4$ is just part 2 of Theorem 2. ■

What remain are the cases where both p and q are even. In these cases we have a new relation. Rotation by π about the x axis, followed by rotation by π about the ℓ axis, equals rotation by $4\pi n/m$ about the z axis:

$$B^{q/2}A^{p/2} = R_\ell^\pi R_x^\pi = R_z^{4\pi n/m} = U^{2n}. \tag{24}$$

To see this, note that $B^{q/2}A^{p/2}$ fixes the z axis while reflecting the x axis across the ℓ axis. The remainder of this paper tracks the effect of this relation.

Theorem 4: *If*

1. *p is even, q is even and m is odd, or*
2. *$\rho(p) \geq 2$, $\rho(q) \geq 2$ and $\rho(m) = 2$, or*
3. *q is even and $\rho(p) \geq \rho(m) \geq 3$,*

   then $G_{n/m}(p,q) = G([p,q],m)$

Proof: We must show that U and $\hat{T}$ are in $G_{n/m}(p,q)$. First we show it suffices to prove that $U^n \in G_{n/m}(p,q)$. Since n and m are relatively prime, U is a power of U^n. And if $U^n \in G_{n/m}(p,q)$ then $R_x^{2\pi/q} = U^nBU^{-n} \in G_{n/m}(p,q)$, and so $\hat{T} \in G_{n/m}(p,q)$. Note that, by eq. (24), we know that U^{2n}, and therefore U^2, is in $G_{n/m}(p,q)$.

If m is odd, then $U^n = (U^{2n})^{(m+1)/2}$, and case 1 is done.

If $\rho(m) = 2$ then $R_z^{\pi/2}$ is an odd power of U, and hence an odd power of U^n. Let s be an odd integer such that $R_z^{\pi/2} = U^{ns}$. $U^{n(s-1)}$ is then an even power of U^n, and hence an element of $G_{n/m}(p,q)$. Thus $R_y^{2\pi/q} = R_z^{-\pi/2}T^pR_z^{\pi/2} = U^{-n(s-1)}BU^{n(s-1)} \in G_{n/m}(p,q)$. Since 4 divided q, a power of this is $R_y^{\pi/2} = S$. Since 4 divides p, $R_x^{\pi/2} = A^{p/4} \in G_{n/m}(p,q)$. Thus $R_z^{\pi/2} = S^{-1}R_x^{\pi/2}S \in G_{n/m}(p,q)$. So $U^n = R_z^{\pi/2}U^{-n(s-1)} \in G_{n/m}(p,q)$, which completes case 2.

In case 3, since $\rho(m) \geq 3$, $R_z^{\pi/2}$ is an even power of U^n and hence is in $G_{n/m}(p,q)$. Since 4 divides p, $R_x^{\pi/2} \in G_{n/m}(p,q)$. From these we see that $S \in G_{n/m}(p,q)$. Conjugating A by S we obtain $R_z^{2\pi/p} \in G_{n/m}(p,q)$. Since $\rho(p) \geq \rho(m)$, $U^b = [R_z^{2\pi/p}]^a \in G_{n/m}(p,q)$, where $a = p/2^{\rho(p)-\rho(m)}$ and $b = m/2^{\rho(m)}$. Since n and b are both odd, $n-b$ is even, so $U^{n-b} = (U^2)^{(n-b)/2} \in G_{n/m}(p,q)$ and $U^n = U^bU^{n-b} \in G_{n/m}(p,q)$. ■

All that is left are a handful of special cases, all of which have p and q even and m divisible by 4. Without loss of generality we assume $\rho(p) \geq \rho(q)$.

Theorem 5:

1. *If $\rho(m) > \rho(p) = \rho(q) = 1$, then*

$$G_{n/m}(p,q) = <\alpha,\beta,\gamma : \alpha^p, \beta^q, \gamma^2, (\alpha\gamma)^2, (\beta\gamma)^2, (\alpha^{p/2}\beta^{q/2})^{m/4}\gamma> . \tag{25}$$

This is a quotient of $D_p *_{\mathbb{Z}_2} D_q$ by the last relation.

2. *If $\rho(m) > \rho(p) > 1$ and $\rho(m) > \rho(q) > 1$, then*

$$G_{n/m}(p,q) = G_{1/2r}(4,4) = G(r,4) *_{D_r} G(r,4), \tag{26}$$

where $r = [p,q,m/2]$ is the least common multiple of p, q and $m/2$, the first $G(r,4)$ subgroup is generated by $R_z^{2\pi/r}$ and $R_x^{\pi/2}$, the second is conjugate to the first by $R_z^{\pi/r}$, and the common D_r subgroup is generated by $R_z^{2\pi/r}$ and R_x^{π}.

3. *If $\rho(m) > \rho(p) > \rho(q) = 1$, then*

$$G_{n/m}(p,q) = G_{1/2s}(4,q) = G(s,4) *_{D_2} D_q, \tag{27}$$

where $s = [p,m/2]$, $G(s,4)$ is generated by $R_z^{2\pi/s}$ and $R_x^{\pi/2}$, D_q is generated by B and R_z^{π}, and the common D_2 subgroup is generated by $B^{q/2}$ and R_z^{π}.

4. *If $\rho(p) > \rho(q) = 1$ and $\rho(m) = 2$, then*

$$G_{n/m}(p,q) = G_{1/2t}(p,2) = D_p *_{D_2} D_t, \tag{28}$$

where $t = [q,m/2]$, D_p is generated by A and R_z^{π}, D_t is generated by $R_z^{2\pi/t}$ and $B^{q/2}$, and the common D_2 is generated by R_z^{π} and $A^{p/2} = R_z^{2\pi/t}B^{q/2}$.

Proof: We begin with case 1. Taking $\alpha = A$, $\beta = B$ and $\gamma = R_z^{\pi} = (A^{p/2}B^{q/2})^{m/4}$ (from (24), since n is odd), it is clear that α, β and γ generate the group (indeed, α and β do), and that all the stated relations hold. We must show that there are no additional relations.

Take an arbitrary word w in α, β and γ and shorten it as follows. Make all exponents of α less than or equal to $p/2$ in magnitude, and all exponents of β less than or equal to $q/2$ in magnitude. Use the relators $(\alpha\gamma)^2$ and $(\beta\gamma)^2$ to move all the γs to the left (i.e., replace $\alpha\gamma$ by $\gamma\alpha^{-1}$ etc.), after which there should be either one or zero γs. If there are any sequences $\cdots\alpha^{p/2}\beta^{q/2}\cdots$ with more than $m/4$ pairs, replace it with γ times a shorter sequence, using the relator $(\alpha^{p/2}\beta^{q/2})^{m/4}\gamma$. Repeat these procedures as necessary until there is at most one γ (at the left), all rotations are by π or less, and there are at most $m/4$ pairs of rotations by π in a row. We claim that this word, if not explicitly the identity, is not equal to the identity.

So consider $w = \gamma^c\alpha^{a_1}\beta^{b_1}\cdots\alpha^{a_n}\beta^{b_n}$, with $c = 0$ or 1 and the aforementioned restrictions on the a_is and b_is. Now rewrite α, β and γ in terms of U and T, getting

$$w = U^{cm/2}\prod_i (T^{qa_i}U^{-n}T^{pb_i}U^n). \tag{29}$$

Since p and q are even, some of the powers of T may be rotations by π, but, since neither p nor q is a multiple of 4, none are rotations by $\pm\pi/2$. We eliminate the $T^{pq/2}$ terms, pushing them to the left (in the same spirit as (23)). If k such pairs are adjacent, then this process changes the $k+1$ flanking $U^{\pm n}$s into a single $U^{\pm(k+1)n}$. However, since k is never bigger than $m/4$, $U^{\pm(k+1)}$ is never a rotation by π, and since n is relatively prime to m neither is $U^{\pm(k+1)n}$. The remaining powers of T are not rotation by multiples of $\pi/2$. Thus, by Extension 2 (applied to U and T), w is nontrivial.

In case 2, since $\rho(p)$ and $\rho(q)$ are at least 2 our group contains rotations by $\pi/2$ about the x and ℓ axes. Since $\rho(m) \geq 3$, our group also contains $R_z^{\pi/2}$ (as in the proof of Theorem 4). Conjugating these rotations about the x and ℓ axes by $R_z^{\pi/2}$ gives rotations by $\pi/2$ about the y and ℓ' axes, where ℓ' is orthogonal to ℓ. Conjugating A and B by these last rotations, we get rotations about the z axis by $2\pi/p$ and $2\pi/q$, and hence by $2\pi/[p,q]$. Since $U^2 = R_z^{4\pi/m}$ is also in the group, and since $\rho(m) > \rho([p,q])$, our group contains $R_z^{2\pi/r}$, where r is the least common multiple $[p,q,m/2]$. Conjugating this by $R_y^{\pi/2}$ and by $R_{\ell'}^{\pi/2}$ gives $R_x^{2\pi/r}$ and $R_\ell^{2\pi/r}$. Finally, since n/m is an odd multiple of $1/2r$, conjugating $R_\ell^{2\pi/r}$ by an appropriate power of $R_z^{2\pi/r}$ gives $R_{\ell''}^{2\pi/r}$, where the axis ℓ'' makes an angle of $2\pi/2r$ with the x-axis. So $G_{1/2r}(r,r) \subseteq G_{n/m}(p,q)$, and therefore $G_{n/m}(p,q) = G_{1/2r}(r,r)$.

Thus $G_{n/m}(p,q)$ does not depend on p, q, n and m separately, but only on r, which is an odd multiple of $m/2$. So we can replace p and q by 4, n by 1 and m by $2r$, which proves the first equality of (26).

To prove the second equality, we again assume that $p = q = 4$, $n = 1$ and $r = m/2$. Our group is now generated by $A = T$ and $B = UTU^{-1}$, where $T = R_x^{\pi/2}$ and $U = R_z^{2\pi/m} = R_z^{\pi/r}$. Our group also contains $U^2 = A^2B^2$. We consider the subgroups generated by U^2 and $A = T$, on the one hand, and by U^2 and B on the other hand. Each group is isomorphic to $G(r,4)$, and they have a common D_r subgroup generated by U^2 and A^2 (or U^2 and $B^2 = A^2U^2$). The group generated by B and A is the quotient of $G(r,4) *_{D_r} G(r,4)$ by any additional relations. We will show that there are no additional relations.

Using (2) and (3) as in the proof of Theorem 1, any element of the first $G(r,4)$ can be written as

$$\alpha = WT^{\pm 1}U^{a_1}T^{\pm 1}\cdots T^{\pm 1}U^{a_N}, \tag{30}$$

where W is an element of D_r, each of the a_i's is even, and, for $i < N$, U^{a_i} is not a rotation by a multiple of $\pi/2$. Similarly, an element of the second $G(r,4)$ can be written as

$$\beta = WUT^{\pm 1}U^{a_1}T^{\pm 1}\cdots T^{\pm 1}U^{a_N-1}. \tag{31}$$

We now need a simple fact about amalgamated free products. Consider some $G *_H \tilde{G}$, and left coset spaces G/H and $\tilde{G}/H$. Pick a set of representatives $\{k_i\}$ (or $\{\tilde{k}_i\}$) for the left cosets of G (or $\tilde{G}$). That is, every element g of G can be written as $g = hk_i$ for some representative k_i and some $h \in H$, with a similar expression for elements of $\tilde{G}$. We claim that any element of $G *_H \tilde{G}$ can be expressed in the form:

$$hg_1\tilde{g}_1g_2\tilde{g}_2g_3\cdots\tilde{g}_p \tag{32}$$

where $h \in H$, each $g_j \in \{k_i\}$ and each $\tilde{g}_j \in \{\tilde{k}_i\}$. To see this, just note that any element of $G *_H \tilde{G}$ can automatically be expressed in the form $\phi_1\tilde{\phi}_1\phi_2\tilde{\phi}_2\phi_3\cdots\tilde{\phi}_p$, with $\phi_j \in G$ and $\tilde{\phi}_j \in \tilde{G}$. Starting from the right, express $\tilde{\phi}_p = \tilde{h}_p \tilde{k}_p$. Then since $\phi_{p-1}\tilde{h}_p \in G$, we can write $h_{p-1}\tilde{k}_{p-1} = \phi_{p-1}\tilde{h}_p$, etc., moving an element of H through the word all the way to the left, where it becomes the h in (32).

By (30), we may choose representatives for the first $G(r,4)/D_r$ of the form $T^{\pm 1}U^{a_1} \cdots T^{\pm 1}U^{a_N}$. Similarly, (31) indicates the form of the representatives for the second $G(r,4)/D_r$. So an arbitrary element of $G(r,4) *_{D_r} G(r,4)$ can be written as

$$\gamma = W(T^{\pm 1}U^{a_{1,1}}T^{\pm 1} \cdots T^{\pm 1}U^{a_{N_1,1}})U(T^{\pm 1}U^{a_{1,2}}T^{\pm 1} \cdots T^{\pm 1}U^{a_{N_2,2}})U^{-1} \cdots, \quad (33)$$

where again W is an arbitrary element of D_r. Notice that none of the powers of U are rotations by multiples of $\pi/2$, except possibly the last term in each parenthetical expression, which then gets multiplied by $U^{\pm 1}$, yielding an odd power of U, hence not a rotation by a multiple of $\pi/2$. Since T is a rotation by $\pi/2$, and none of the powers of U are, Extension 2 implies the resulting word is not equal to the identity, which completes case 2.

In case 3, as in case 2, we have rotations by $\pi/2$ about the x and z axes, hence around the y axis, and hence can transfer generators from the x to the z axis and vice-versa. Defining $U = R_z^{\pi/s}$, as in case 2 we again have U^2 in our group, and, by conjugating by a power of U^2, we can exchange the ℓ axis for an axis that makes an angle of π/s with the x axis. This shows that p and m do not contribute separately, but only through their least common multiple $2s$, and that n does not matter at all. This establishes the first equality in (27).

For the second equality we assume $p = 4$ and $n = 1$, so $m = 2s$ and $U = R_z^{2\pi/m}$. The $G(s,4)$ and D_q subgroups are manifest, as is their common D_2 subgroup. All that remains is to show that there are no hidden relations among the elements of $G(s,4) *_{D_2} D_q$.

The rest of the argument goes just as in case 2. Find representatives for all the left cosets of D_2 in $G(s,4)$ and in D_q, write out an arbitrary element of the amalgamated free product, and notice that it isn't trivial.

For case 4, we define $U = R_z^{2\pi/m}$. Since $\rho(m) = 2$, $R_z^{\pi/2}$ is an odd power of U. Since n is odd, conjugating B by an even power of U gives a rotation about the y axis, and conjugating this by $A^{p/4}$ gives a rotation about the z axis. Thus generators may be transfered back and forth between the ℓ and z axes, demonstrating that q and m only contribute through their least common multiple $2t$. Also, the result is independent of n as usual. This gives the first equality.

For the second equality, assume $q = 2$, so $t = m/2$. The same argument as in cases 2 and 3 again works. ■

§4. Conclusions and Remarks

The group of rotations of three dimensional Euclidean space is a fundamental object in mathematics and in mathematical models of science. We have described above how a few elementary facts about rotations by $\pi/2$ determine the relations between a wide range of pairs of rotations: those of finite order, about axes that are themselves separated by an angle of finite order.

Two-generator subgroups of $SO(3)$ have been studied in the context of the Banach-Tarski paradox (see [W]). There one uses a pair of "independent" rotations, that is, rotations which generate a free subgroup of $SO(3)$, to produce interesting subsets of the unit sphere. Since in a free group there are NO relations, such rotations must have infinite order, being rotations by irrational angles.

At first glance such rotations would seem different, indeed complementary, to the subject of this paper. However, it is not difficult to exhibit explicit free 2-generator subgroups of many of the groups considered here. For example, let m be any positive integer other than 1, 2, 4 or 8, let $S = R_y^{\pi/2}$, and let $T = R_x^{2\pi/m}$. Then the two rotations $A = STST$ and $B = ST^2ST^2$ are independent. To see this, write an arbitrary word in A and B in terms of S and T. Since neither T nor T^2 is a rotation by a multiple of $\pi/2$, the result is a word in S and T of the form (5), and is therefore not the identity. (It is not difficult to check that the word in S and T is always longer than the original word in A and B.) The group generated by A and B is then a free 2-generator subgroup of $\hat{G}(m, 4)$.

Aknowledgements. We thank John Conway, J.-P. Serre and John Tate for helpful discussions. This work, as well as [RS], is an outgrowth of work with John Conway [CR]. We also thank Stuart Levy for writing the graphics software from which Figure 1 was obtained.

References

[CR] J. Conway and C. Radin: Quaquaversal tilings and rotations. Preprint, Princeton University and University of Texas (1995) [Obtainable from the electronic archive: mp_arc@math.utexas.edu]

[RS] C. Radin and L. Sadun: Subgroups of $SO(3)$ associated wtih tilings. Preprint, University of Texas (1996) [Obtainable from the electronic archive: mp_arc@math.utexas.edu]

[W] S. Wagon: The Banach-Tarski Paradox. Cambridge: The University Press 1985



Figure 1. Part of a quaquaversal tiling


$S = R_y^{\pi/2}$
$B = R_\ell^{2\pi/q}$
Y
ℓ
$A = R_x^{2\pi/p}$
$T, \hat{T} = R_x^{\text{various angles}}$
X
Z
$U = R_z^{2\pi/m}$
$C = R_z^{2\pi/q}$

Figure 2